

NAME (Printed): \_\_\_\_\_

Math 304-6      Linear Algebra      Fall 2025      Quiz 4      Feingold

INSTRUCTIONS: Show all calculations needed to justify your answers.

- (1) (5 Pts) Let  $m, n, p$  be given positive integers and let  $A \in \mathbb{F}_n^m$  be a given  $m \times n$  matrix fixed for this problem. Then  $A$  determines a set of  $n \times p$  matrices  $W = \{B \in \mathbb{F}_p^n \mid AB = 0_p^m\}$ . Show  $W$  is a subspace of  $\mathbb{F}_p^n$ .

- 
- (2) (5 Pts) Let  $A = \begin{bmatrix} 1 & 3 \\ 3 & 9 \end{bmatrix} \in \mathbb{F}_2^2$ . Find  $W = \left\{ B = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{F}_2^2 \mid AB = 0_2^2 \right\}$  by solving a linear system.

INSTRUCTIONS: Show all calculations needed to justify your answers.

- (1) (5 Pts) Let  $m, n, p$  be given positive integers and let  $A \in \mathbb{F}_n^m$  be a given  $m \times n$  matrix fixed for this problem. Then  $A$  determines a set of  $n \times p$  matrices  $W = \{B \in \mathbb{F}_p^n \mid AB = 0_p^m\}$ . Show  $W$  is a subspace of  $\mathbb{F}_p^n$ .

SOLUTION: First show  $W$  is closed under addition. Let  $B_1, B_2 \in W$ , so  $AB_1 = 0_p^m$  and  $AB_2 = 0_p^m$ . Then  $A(B_1 + B_2) = AB_1 + AB_2 = 0_p^m + 0_p^m = 0_p^m$  means  $B_1 + B_2 \in W$ .

Next show  $W$  is closed under scalar multiplication. Let  $B \in W$  so  $AB = 0_p^m$ , and let  $r \in \mathbb{F}$ . Then  $A(rB) = r(AB) = r0_p^m = 0_p^m$  means  $rB \in W$ .

Finally,  $0_p^n \in W$  since  $A0_p^n = 0_p^m$ .

- (2) (5 Pts) Let  $A = \begin{bmatrix} 1 & 3 \\ 3 & 9 \end{bmatrix} \in \mathbb{F}_2^2$ . Find  $W = \left\{ B = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{F}_2^2 \mid AB = 0_2^2 \right\}$  by solving a linear system.

SOLUTION: To find all  $B \in \mathbb{F}_2^2$  such that  $AB = 0_2^2$ , write out the condition as

$$\begin{bmatrix} 1 & 3 \\ 3 & 9 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} (a+3c) & (b+3d) \\ (3a+9c) & (3b+9d) \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

This gives us four equations in four variables which we solve by row reduction as follows:

$$\text{Row reduce } \left[ \begin{array}{cccc|c} 1 & 0 & 3 & 0 & 0 \\ 0 & 1 & 0 & 3 & 0 \\ 3 & 0 & 9 & 0 & 0 \\ 0 & 3 & 0 & 9 & 0 \end{array} \right] \text{ to } \left[ \begin{array}{cccc|c} 1 & 0 & 3 & 0 & 0 \\ 0 & 1 & 0 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right] \text{ so } \begin{array}{l} a = -3c \\ b = -3d \\ c \in \mathbb{F} \text{ free} \\ d \in \mathbb{F} \text{ free} \end{array}$$

$$\text{so } W = \left\{ B = \begin{bmatrix} -3c & -3d \\ c & d \end{bmatrix} \in \mathbb{F}_2^2 \mid c, d \in \mathbb{F} \right\}.$$