

NAME (Printed): \_\_\_\_\_

Math 404    Linear Algebra    Spring 2023    Quiz 1    Feingold

**Show all calculations needed to justify your answers.**

Let  $L : \mathbf{F}_2^2 \rightarrow \mathbf{F}_2^2$  be the function  $L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & b - c \\ 0 & 0 \end{bmatrix}$ .

(1) (1 pt) For  $A_1 = \begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix}$  and  $A_2 = \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix}$ , show  $L(A_1 + A_2) = L(A_1) + L(A_2)$ .

---

(2) (1 pt) For  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  and  $r \in \mathbf{F}$ , show that  $L(rA) = rL(A)$ .

---

(3) (2 pts) Since parts (1) and (2) show that  $L$  is linear, find the subspace  $\text{Ker}(L)$ .

---

(4) (1 pt) How does part (3) show that the set of all **symmetric** matrices in  $\mathbf{F}_2^2$  is a subspace?

Show all calculations needed to justify your answers.

Let  $L : \mathbf{F}_2^2 \rightarrow \mathbf{F}_2^2$  be the function  $L \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} 0 & b - c \\ 0 & 0 \end{bmatrix}$ .

(1) (1 pt) For  $A_1 = \begin{bmatrix} a_1 & b_1 \\ c_1 & d_1 \end{bmatrix}$  and  $A_2 = \begin{bmatrix} a_2 & b_2 \\ c_2 & d_2 \end{bmatrix}$ , show  $L(A_1 + A_2) = L(A_1) + L(A_2)$ .

**Solution:**  $A_1 + A_2 = \begin{bmatrix} (a_1 + a_2) & (b_1 + b_2) \\ (c_1 + c_2) & (d_1 + d_2) \end{bmatrix}$  so

$$L(A_1 + A_2) = \begin{bmatrix} 0 & (b_1 + b_2) - (c_1 + c_2) \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & (b_1 - c_1) \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} 0 & (b_2 - c_2) \\ 0 & 0 \end{bmatrix} = L(A_1) + L(A_2).$$

(2) (1 pt) For  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  and  $r \in \mathbf{F}$ , show that  $L(rA) = rL(A)$ .

**Solution:**  $L(rA) = L \begin{bmatrix} ra & rb \\ rc & rd \end{bmatrix} = \begin{bmatrix} 0 & rb - rc \\ 0 & 0 \end{bmatrix} = r \begin{bmatrix} 0 & b - c \\ 0 & 0 \end{bmatrix} = rL(A)$ .

(3) (2 pts) Since parts (1) and (2) show that  $L$  is linear, find the subspace  $\text{Ker}(L)$ .

**Solution:**

$$\text{Ker}(L) = \left\{ A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{F}_2^2 \mid L(A) = \begin{bmatrix} 0 & b - c \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \right\} =$$

$$\left\{ A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{F}_2^2 \mid b = c \right\} = \left\{ A = \begin{bmatrix} a & b \\ b & d \end{bmatrix} \in \mathbf{F}_2^2 \mid a, b, d \in \mathbf{F} \right\} = \{A \in \mathbf{F}_2^2 \mid A^T = A\}.$$

(4) (1 pt) How does part (3) show that the set of all **symmetric** matrices in  $\mathbf{F}_2^2$  is a subspace?

**Solution:** In part (3) we found that  $\text{Ker}(L)$  is the set of all **symmetric** matrices in  $\mathbf{F}_2^2$ , so as a kernel it is a subspace.