

Homework

due on Monday, April 25

Read carefully chapter 11 of Joseph's book and chapters 5, 6 of Dunham's book. Read the article by N. Schappacher on Diophantus (linked on the course web page). Solve the following problems.

Problem 1. a) Consider a convex quadrilateral $ABCD$ inscribed in a circle with center O such that AD is the diameter, $\angle AOB = 2y$, $\angle AOC = 2x$, $\angle COD = 2z$. Explain how to use Ptolemy's theorem to prove that

$$\sin(x - y) = \sin x \cos y - \sin y \cos x$$

and

$$\cos(y + z) = \cos y \cos z - \sin y \sin z.$$

b) Consider a convex quadrilateral $ABCD$ inscribed in a circle with center O such that AC is the diameter, $\angle AOB = 2y$, $\angle AOD = 2x$, $\angle COB = 2z$. Explain how to use Ptolemy's theorem to prove that

$$\sin(x + y) = \sin x \cos y + \sin y \cos x$$

and

$$\cos(z - x) = \cos z \cos x + \sin z \sin x$$

(assuming $z > x$).

Problem 2. a) What is the ratio of the diagonal in a regular pentagon to its side? (Euclid, Book XIII, Prop. 8). Explain how this implies that

$$\sin 72^\circ = \frac{\sqrt{5} + 1}{2} \sin 36^\circ.$$

Use this equality and the formula for $\sin 2x$ to show that $\cos 36^\circ = (\sqrt{5} + 1)/4$. Conclude that $\sin 36^\circ = \sqrt{10 - 2\sqrt{5}}/4$.

b) Use a) to prove that $\sin 18^\circ = (\sqrt{5} - 1)/4$ and $\cos 18^\circ = \sqrt{10 + 2\sqrt{5}}/4$. Show that $\sin 15^\circ = (\sqrt{6} - \sqrt{2})/4$ and $\cos 15^\circ = (\sqrt{6} + \sqrt{2})/4$ (hint: $15 = 45 - 30$). Then show that

$$\sin 3^\circ = \frac{\sqrt{30} + \sqrt{10} - \sqrt{6} - \sqrt{2} + 2\sqrt{5 + \sqrt{5}} - 2\sqrt{15 + 3\sqrt{5}}}{16}$$

and

$$\cos 3^\circ = \frac{\sqrt{30} - \sqrt{10} - \sqrt{6} + \sqrt{2} + 2\sqrt{5} + \sqrt{5} + 2\sqrt{15 + 3\sqrt{5}}}{16}.$$

Hint: $3 = 18 - 15$, and $\cos(x - y) = \cos x \cos y + \sin x \sin y$.

Problem 3. Learn a proof of the Theorem of Menelaus (for example, from the links provided on the course web page). Use this theorem to prove the following results.

a) The lines AA_1 , BB_1 and CC_1 intersect in one point O . Let the lines AB and A_1B_1 intersect at C_2 , the lines AC and A_1C_1 intersect at B_2 , and the lines BC and B_1C_1 intersect at A_2 . Prove that the points A_2 , B_2 , C_2 are collinear. This is often called Desargues Theorem. Hint. Apply Menelaus' Theorem to triangles OAB , OBC , OAC and appropriate lines. Then apply its converse to the triangle ABC .

b) Points A_1 , B_1 , C_1 are collinear and so are points A_2 , B_2 , C_2 . The lines A_1B_2 and A_2B_1 intersect at a point C , the lines A_1C_2 and A_2C_1 intersect at a point B , and the lines B_1C_2 and B_2C_1 intersect at a point A . Prove that the points A , B , C are collinear. This is Pappus' Theorem. Hint. Let A_0 , B_0 , C_0 be the vertices of the triangle determined by the lines A_1B_2 , B_1C_2 , and C_1A_2 (where A_0 is the point of intersection of A_1B_2 and A_2C_1 , etc.). Apply Menelaus' Theorem to the triangle $A_0B_0C_0$ and five appropriate lines.

Problem 4. Justify Newton's solution of the duplication of a cube, as described in the link on the course web page, by following the steps below (the notation is from the link). Alternatively, find your own justification.

a) Let $AG = x$, $CG = y$. Show that $\angle ACG = \pi/2$ and derive a relation between x and y .

b) Show that $DC = \sqrt{3}$.

c) Apply the Theorem of Menelaus to the triangle ADG . This should give you another relation between x and y .

d) Find x and y .

Problem 5. Let $ABCD$ be a convex quadrilateral. Let F be the point of intersection of its diagonals AC and BD . Set $a = AB$, $b = BC$, $c = CD$, $d = DA$, $p = (a + b + c + d)/2$, $e = AC$, $f = BD$, $\alpha = \angle AFB$.

a) Prove that $a^2 + c^2 - b^2 - d^2 = -2ef \cos \alpha$. Hint: Apply the law of cosines to triangles AFB , BFC , CFD , DFA .

b) Prove that the area S of the quadrilateral is given by $2S = ef \sin \alpha$. Hint: Add areas of the triangles AFB , BFC , CFD , DFA .

c) Show that $4(p-a)(p-c) = b^2 + d^2 - a^2 - c^2 + 2(ac + bd)$ and $4(p-b)(p-d) = a^2 + c^2 - b^2 - d^2 + 2(ac + bd)$. Conclude that $16(p-a)(p-b)(p-c)(p-d) = 4[(ac + bd)^2 - (ef)^2] + 16S^2$. Hint: Use a) and b).

d) Prove that if $ABCD$ is cyclic then $S = \sqrt{(p-a)(p-b)(p-c)(p-d)}$. Hint: Use Ptolemy's Theorem.

The formula in d) generalizes Heron's Theorem. It has been stated by an Indian mathematician and astronomer Brahmagupta (598-670 AD), but we do not know how he derived this formula.

Remark: One can prove that $(ac)^2 + (bd)^2 - (ef)^2 = 2abcd \cos 2\theta$, where $2\theta = \angle DAB + \angle BCD$. Using this one can show that $S^2 = (p-a)(p-b)(p-c)(p-d) - abcd \cos^2 \theta$.