

## Homework 1

due on Monday, February 14

Read carefully Chapter 1 of Miln's book and sections 1.1-1.6, 2.1, 2.3, 2.4, 2.5, 3.1, 3.2, 3.3 of Dummit and Foote.

**Problem 1.** Let  $G$  be a group. Recall that  $(m, n)$  is the greatest common divisor of  $m$  and  $n$ . Prove that:

- a) If  $a \in G$  has finite order  $n$  then, for any integer  $k$ , the order of  $a^k$  is  $n/(n, k)$ .
- b) If  $a$  has order  $m$ ,  $b$  has order  $n$ , and  $ab = ba$  then the order of  $ab$  divides  $mn/(m, n)$  and is divisible by  $mn/(m, n)^2$ .
- c) If  $G$  has an element  $a$  of order  $m$  and an element  $b$  of order  $n$  such that  $ab = ba$  then  $G$  has an element of order  $[m, n]$  ( $[m, n]$  is the least common multiple of  $m$  and  $n$ ).
- d) If  $G$  is a finite abelian group and  $N$  is the smallest positive integer such that  $g^N = e$  for all  $g \in G$ , then  $G$  has an element of order  $N$ .

**Remark.** In general, the  $N$  defined in d) makes sense for any group (it can be infinite) and it is called **the exponent of  $G$** .

- e) If  $f : G \rightarrow H$  is a homomorphism and  $a \in G$  has finite order  $n$ , then  $f(a)$  has also finite order  $k$  which divides  $n$ . Also,  $a^m$  is in the kernel of  $f$  iff  $k$  divides  $m$ .
- f) Let  $G$  be a cyclic group of order  $n$  and  $H$  a cyclic group of order  $m$  (we allow the orders to be infinite). Show that the set of all homomorphism from  $G$  to  $H$  is a group with operation  $+$  defined by  $(f + g)(a) = f(a)g(a)$  (this is true for arbitrary  $G$  and abelian  $H$ ). Describe this group for each pair  $m, n$ .

- g) Study Theorem 1.64 and its proof in Miln's book.

**Problem2.** Let  $G$  be a group and  $H$  its subgroup.

- a) Show that if  $a_i H$ ,  $i \in I$  are all the left cosets of  $H$  in  $G$  then  $Ha_i^{-1}$ ,  $i \in I$  are all the right cosets of  $H$  in  $G$  (each listed once). Conclude that the number of left cosets of  $H$  is finite iff the number of right cosets is finite and these numbers coincide. The number of left (right) cosets of  $H$  in  $G$  is called the **index** of  $H$  in  $G$  and it is usually denoted by  $[G : H]$ .
- b) Prove that if  $K < H < G$  then  $[G : K] = [G : H][H : K]$ .
- c) Show that for any subgroup  $K$  of  $G$  we have  $[K : H \cap K] \leq [G : H]$ .
- d) Prove that if  $H, K$  are subgroups of  $G$  of finite index then so is  $H \cap K$  and  $[G : H \cap K] \leq [G : H][G : K]$ .
- e) Prove that if  $H$  is of finite index then  $G$  is finitely generated iff  $H$  is finitely generated.
- f) Prove that if  $H$  is of finite index then there is a normal subgroup of  $G$  of finite index contained in  $H$  (show that the number of conjugates of  $H$  is finite and take their intersection).
- g) Show that if  $G$  is finitely generated then it has only finitely many subgroups of a given finite index  $n$  (use the fact that the action of  $G$  on cosets of a subgroup  $K$  of index  $n$  defines a homomorphism of  $G$  into  $S_n$  whose kernel is contained in  $K$ ).

h) If  $[G : H] = 2$  then  $H$  is normal.

i) Show that if  $[G : H] = n$  then  $g^{n!} \in H$  for all  $g \in G$ . If  $H$  is normal then  $n!$  can be replaced by  $n$ . Show that without normality this is no longer true.

**Problem 3.** Let  $G$  be the set of all bijections  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  which preserve distance, i.e. such that  $|f(i) - f(j)| = |i - j|$  for all integers  $i, j$ .

a) Show that  $G$  is a subgroup of  $\text{Sym}(\mathbb{Z})$ . It is called the **infinite dihedral group** and it is often denoted by  $D_\infty$ .

b) The group  $G$  contains elements  $T, S$  such that  $T(a) = a + 1$  and  $S(a) = -a$  for all integers  $a$ . Prove that  $S * T = T^{-1} * S$ . Show that the subgroup  $\langle T \rangle$  is infinite. What is  $\langle S \rangle$ ?

c) Show that if  $F \in G$  and  $F(0) = 0$  then either  $F = 1$  (the identity) or  $F = S$ .

d) Show that every element of  $G$  is of the form  $T^i$  or  $ST^i$  for some integer  $i$  (try to use similar argument to the one we used for dihedral group of order  $n$ ).

e) Suppose that  $T^5 S^7 T^3 = S^a T^b$ . Find  $a$  and  $b$ .

f) Find the center and the derived subgroup of  $G$ .

**Problem 4.** a) Describe all subgroups and normal subgroups of  $D_n$ .

b) Describe the center and the derived group of  $D_n$ .

c) For which  $m, n$  is there a surjective homomorphism from  $D_m$  to  $D_n$ ? (**Optional:** Describe all homomorphisms from  $D_m$  to  $D_n$ .)

d) Prove that if  $x, y$  are two elements of order 2 in a group  $G$  and  $xy \neq yx$  then the subgroup  $\langle x, y \rangle$  of  $G$  is isomorphic to a dihedral group (finite or infinite).

**Problem 5.** In the group  $GL_2(\mathbb{C})$  of all invertible  $2 \times 2$  matrices with entries in complex numbers consider the matrices  $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ ,  $i = \begin{pmatrix} \sqrt{-1} & 0 \\ 0 & -\sqrt{-1} \end{pmatrix}$ ,  $j = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ ,  $k = ij = \begin{pmatrix} 0 & \sqrt{-1} \\ \sqrt{-1} & 0 \end{pmatrix}$ . Let  $Q_8$  be the set  $\{I, -I, i, -i, j, -j, k, -k\}$ .

a) Show that  $Q_8$  is a subgroup of  $GL_2(\mathbb{C})$ . Write the table of multiplication in  $Q_8$ .  $Q_8$  is called the **quaternion group**.

b) List all subgroups of  $Q_8$ .

Furthermore, solve problems 18, 23 to 1.6, problem 6 to 2.1, problem 26 to 2.3.