

1. δ -FUNCTORS

Let \mathcal{C}, \mathcal{D} be abelian categories. Recall that for any abelian category \mathcal{A} we have the category $SES(\mathcal{A})$ of short exact sequences in \mathcal{A} and three functors $T_i : SES(\mathcal{A}) \longrightarrow \mathcal{A}$, $i = 1, 2, 3$, defined by

$$T_i(0 \longrightarrow A_1 \longrightarrow A_2 \longrightarrow A_3 \longrightarrow 0) = A_i.$$

Inspired by the results of the previous section we make the following definitions

Definition 1. A δ -functor from \mathcal{C} to \mathcal{D} is a collection of additive functors $F^n : \mathcal{C} \longrightarrow \mathcal{D}$, $n \in \mathbb{Z}$, and natural transformations $\delta_n : F^n \circ T_3 \longrightarrow F^{n+1} \circ T_1$ of functors from $SES(\mathcal{C})$ to \mathcal{D} such that for every short exact sequence $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$ in \mathcal{C} the sequence

$$\dots \longrightarrow F^{n-1}(C) \xrightarrow{\delta_{n-1}} F^n(A) \longrightarrow F^n(B) \longrightarrow F^n(C) \xrightarrow{\delta_n} F^{n+1}(A) \longrightarrow \dots$$

is a complex. If this complex is exact for every short exact sequence in \mathcal{C} then the δ -functor is called **exact**.

Definition 2. A δ -functor is called **cohomological (homological)** if $F^n = 0$ for all $n < 0$ (resp. for all $n > 0$).

Remark. For a homological δ -functor it is customary to write F_n for F^{-n} , etc.

Definition 3. A cohomological δ -functor is called **effaceable** if for any object X of \mathcal{C} there is a monomorphism $u : X \longrightarrow Y$ such that $F^n(u) = 0$ for all $n > 0$. A homological δ -functor is called **effaceable** if for any

object X of \mathcal{C} there is an epimorphism $u : Y \longrightarrow X$ such that $F^n(u) = 0$ for all $n < 0$. Any such u is called an **effacing** morphism.

Definition 4. A cohomological (or homological) δ -functor is called **universal**, if it is exact and effaceable.

Theorem 1. Let (F^n) be a universal cohomological (homological) δ -functor and (G^n) any δ -functor. Any natural transformation $\alpha : F^0 \longrightarrow G^0$ of functors (resp. $\alpha : G^0 \longrightarrow F^0$) uniquely extends to a natural transformation of δ -functors.

Remark. A natural transformation of δ -functors is a collection of natural transformations $\alpha_n : F^n \longrightarrow G^n$, $n \in \mathbb{Z}$, such that for every k and every short exact sequence $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$ in \mathcal{C} the following diagram commutes:

$$(1) \quad \begin{array}{ccc} F^{k-1}(C) & \xrightarrow{\delta_{k-1}} & F^k(A) \\ \downarrow \alpha_{k-1} & & \downarrow \alpha_k \\ G^{k-1}(C) & \xrightarrow{\delta_{k-1}} & G^k(A) \end{array}$$

Proof: We give a proof for cohomological δ -functors and leave the case of homological δ -functors as an exercise.

Uniqueness. Suppose we have two natural transformations $(\alpha_n), (\beta_n)$ such that $\alpha_0 = \alpha = \beta_0$. Clearly $\alpha_n = 0 = \beta_n$ for all $n < 0$, since $F^n = 0$ for such n . We show that $\alpha_n = \beta_n$ by induction on n . For $n = 0$ it is our assumption. Suppose then that $\alpha_k = \beta_k$ for all $k < n$. Let X be an object of \mathcal{C} and $u : X \longrightarrow Y$ an effacing monomorphism. Thus we have an exact sequence

$$0 \longrightarrow X \xrightarrow{u} Y \xrightarrow{j} Z \longrightarrow 0$$

where $Z = \text{coker } u$, which leads to the following commutative diagram

$$\begin{array}{ccccc}
F^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & F^n(X) & \xrightarrow{F^n(u)} & F^n(Y) \\
\alpha_{n-1}=\beta_{n-1} \downarrow & & \alpha_n \downarrow \beta_n & & \\
G^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & G^n(X) & \xrightarrow{G^n(u)} & G^n(Y)
\end{array}$$

with the top row exact. Note that $\alpha_n \delta_{n-1} = \delta_{n-1} \alpha_{n-1} = \delta_{n-1} \beta_{n-1} = \beta_n \delta_{n-1}$. Since $F^n(u) = 0$, the morphism δ_{n-1} in the top row is an epimorphism, hence the equality $\alpha_n \delta_{n-1} = \beta_n \delta_{n-1}$ implies that $\alpha_n = \beta_n$.

Existence. Suppose that we have already constructed natural transformations α_k for $k < n$ such that the diagrams (1) commute for $k < n$. Let X be an object of \mathcal{C} and $u : X \rightarrow Y$ an effacing monomorphism. Thus we have an exact sequence

$$0 \rightarrow X \xrightarrow{u} Y \xrightarrow{j} Z \rightarrow 0$$

where $Z = \text{coker } u$, which leads to the following commutative diagram

$$\begin{array}{ccccccc}
F^{n-1}(Y) & \xrightarrow{F^{n-1}(j)} & F^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & F^n(X) & \longrightarrow & 0 \\
\alpha_{n-1} \downarrow & & \alpha_{n-1} \downarrow & & & & \\
G^{n-1}(Y) & \xrightarrow{G^{n-1}(j)} & G^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & G^n(X) & &
\end{array}$$

with the top row exact. Note that $(\delta_{n-1} \alpha_{n-1}) F^{n-1}(j) = \alpha_{n-1} G^{n-1}(j) \delta_{n-1} = 0$ (since $G^{n-1}(j) \delta_{n-1} = 0$). It follows that there exists unique morphism α^u such that the diagram

$$\begin{array}{ccc}
F^{n-1}(K) & \xrightarrow{\delta_{n-1}} & F^n(X) \\
\downarrow \alpha_{n-1} & & \downarrow \alpha^u \\
G^{n-1}(K) & \xrightarrow{\delta_{n-1}} & G^n(X)
\end{array}$$

commutes. If the natural transformation α_n exists, then we must have $\alpha_n = \alpha^u$ by uniqueness. In particular, α^u should not depend on the effacing morphism u . We will show that this is indeed true and that setting $\alpha_n = \alpha^u$ indeed defines a natural transformation of functors for which the diagrams (1) commute for $k = n$.

Suppose that we are given a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & X & \xrightarrow{u} & Y & \xrightarrow{j} & Z \longrightarrow 0 \\ & & \downarrow f & & \downarrow & & \downarrow g \\ 0 & \longrightarrow & A & \xrightarrow{w} & B & \xrightarrow{i} & C \longrightarrow 0 \end{array}$$

with exact rows and u, w effacing monomorphisms. This leads to the following cube

$$\begin{array}{ccccc} F^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & F^n(X) & & \\ \downarrow F^{n-1}(g) & \searrow \alpha_{n-1} & \downarrow F^n(f) & \searrow \alpha^u & \\ & G^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & G^n(X) & \\ & \downarrow G^{n-1}(g) & & \downarrow G^n(f) & \\ F^{n-1}(C) & \xrightarrow{\delta_{n-1}} & F^n(A) & & \\ & \searrow \alpha_{n-1} & \searrow \alpha^w & & \\ & G^{n-1}(C) & \xrightarrow{\delta_{n-1}} & G^n(A) & \end{array}$$

in which all faces except perhaps the right face commute and in which the top δ_{n-1} is an epimorphism. We claim that it follows that the right face commutes too, i.e. that $G^n(f)\alpha^u = \alpha^w F^n(f)$. In fact, since the top δ_{n-1} is an epimorphism, it is enough to show that $G^n(f)\alpha^u\delta_{n-1} = \alpha^w F^n(f)\delta_{n-1}$. The commutativity of the faces gives

$$\begin{aligned} G^n(f)\alpha^u\delta_{n-1} &= G^n(f)\delta_{n-1}\alpha_{n-1} = \delta_{n-1}G^{n-1}(g)\alpha_{n-1} = \\ &= \delta_{n-1}\alpha_{n-1}F^{n-1}(g) = \alpha^w\delta_{n-1}F^{n-1}(g) = \alpha^w F^n(f)\delta_{n-1}, \end{aligned}$$

which confirms our claim.

Suppose now that we have two short exact sequences

$$0 \longrightarrow X \xrightarrow{u} Y \xrightarrow{j} Z \longrightarrow 0$$

and

$$0 \longrightarrow A \xrightarrow{w} B \xrightarrow{i} C \longrightarrow 0$$

with u, w effacing monomorphisms and a morphism $f : X \longrightarrow A$. We can not apply the previous paragraph directly since there is no reason in general for f to extend to a morphism of short exact sequences. Fortunately, there is a way around this problem. We construct a third exact sequence:

$$0 \longrightarrow X \xrightarrow{u \times wf} A \times Y \longrightarrow W \longrightarrow 0$$

(where W is a cokernel of $u \times wf$). The following diagram commutes

$$\begin{array}{ccccccccc} 0 & \longrightarrow & X & \xrightarrow{u} & Y & \xrightarrow{j} & Z & \longrightarrow & 0 \\ & & \uparrow id & & \uparrow p_Y & & \uparrow & & \\ 0 & \longrightarrow & X & \xrightarrow{u \times wf} & Y \times B & \longrightarrow & W & \longrightarrow & 0 \\ & & \downarrow f & & \downarrow p_B & & \downarrow & & \\ 0 & \longrightarrow & A & \xrightarrow{w} & B & \xrightarrow{i} & C & \longrightarrow & 0 \end{array}$$

where p_B, p_Y are projections and the vertical arrows from W are uniquely determined by the universal property of a cokernel. Since F^n are additive, $F^n(u \times wf) = F^n(u) \times F^n(w)F^n(f) = 0 \times 0 = 0$ for $n > 0$. Thus $u \times wf$ is an effacing monomorphism. Applying the previous paragraph to the top two and bottom two exact sequences yields the equalities $G^n(id)\alpha^{u \times wf} = \alpha^u F^n(id)$ and $G^n(f)\alpha^{u \times wf} = \alpha^w F^n(f)$. Thus $\alpha^{u \times wf} = \alpha^u$ and $G^n(f)\alpha^u = \alpha^w F^n(f)$. In particular, we may take $A = X$ and $f = id$ and we see that $\alpha^u = \alpha^w$. This proves independence of the effacing monomorphism and allows us to define $\alpha_n = \alpha^u$. Since for any morphism f we have $G^n(f)\alpha_n = \alpha_n F^n(f)$, we see that α_n is a natural transformation from F^n to G^n .

It remains to show that the diagrams (1) commute for $k = n$. So we start with an exact sequence $0 \longrightarrow A \xrightarrow{i} B \longrightarrow C \longrightarrow 0$ in \mathcal{C} . Note that if i is an effacing monomorphism then the corresponding diagram (1) commutes for $k = n$ by definition of α_n . Let $u : B \longrightarrow Y$ be an effacing monomorphism. Note that $ui : A \longrightarrow Y$ is also an effacing

monomorphism. We have a commutative diagram

$$\begin{array}{ccccccc}
 0 & \longrightarrow & A & \xrightarrow{i} & B & \longrightarrow & C \longrightarrow 0 \\
 & & \downarrow id & & \downarrow u & & \downarrow w \\
 0 & \longrightarrow & A & \xrightarrow{ui} & Y & \longrightarrow & Z \longrightarrow 0
 \end{array}$$

which leads to the following cube

$$\begin{array}{ccccc}
 F^{n-1}(C) & \xrightarrow{\delta_{n-1}} & F^n(A) & & \\
 \downarrow F^{n-1}(w) & \searrow \alpha_{n-1} & \downarrow F^n(id) & \searrow \alpha_n & \\
 & G^{n-1}(C) & \xrightarrow{\delta_{n-1}} & G^n(A) & \\
 & \downarrow & & \downarrow G^n(id) & \\
 F^{n-1}(Z) & \xrightarrow{G^{n-1}(w)} & F^n(A) & & \\
 \searrow \alpha_{n-1} & \downarrow & \searrow \alpha_n & & \\
 & G^{n-1}(Z) & \xrightarrow{\delta_{n-1}} & G^n(A) &
 \end{array}$$

in which all faces except possibly the top one commute. Our proof will be completed if we show that the top face commutes as well. But this is quite simple at this point:

$$\begin{aligned}
 \alpha_n \delta_{n-1} &= G^n(id) \alpha_n \delta_{n-1} = \alpha_n F^n(id) \delta_{n-1} = \alpha_n \delta_{n-1} F^{n-1}(w) = \\
 &= \delta_{n-1} \alpha_{n-1} F^{n-1}(w) = \delta_{n-1} G^{n-1}(w) \alpha_{n-1} = G^n(id) \delta_{n-1} \alpha_{n-1} = \delta_{n-1} \alpha_{n-1}. \quad \square
 \end{aligned}$$

Corollary 1. *A universal cohomological (or homological) δ -functor (F^n) is uniquely determined by F^0 . More precisely, if (G^n) is another universal cohomological (or homological) δ -functor and $\alpha : F^0 \longrightarrow G^0$ (resp. $\alpha : G^0 \longrightarrow F^0$) is an equivalence of functors then it uniquely extends to an equivalence (α_n) of δ -functors.*

Proof: Let β be the inverse of α . There are unique extensions (α_n) of α and (β_n) of β , as given by Theorem 1. Now $\gamma_n = \alpha_n \beta_n$ is an extension of $\alpha \beta = id$. Since identity $\gamma'_n = id$ is also an extension, the uniqueness in

Theorem 1 implies that $\gamma_n = \alpha_n \beta_n = id$ for all n . Similarly, $\beta_n \alpha_n = id$ for all n . \square

The last lemma indicates that a universal δ -functor can be recovered from F^0 . It is natural to ask for a way to compute the δ -functor from F^0 . A related question is what functors $F : \mathcal{C} \longrightarrow \mathcal{D}$ are equal to F^0 for some universal δ -functor. The following simple observation gives a necessary condition:

Proposition 1. *If (F^n) is a universal cohomological (homological) δ -functor then F^0 is left exact (resp. right exact).*

Proof: If $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$ is a short exact sequence in \mathcal{C} then we have the long exact sequence

$$\dots \longrightarrow F^{-1}(C) \longrightarrow F^0(A) \longrightarrow F^0(B) \longrightarrow F^0(C) \longrightarrow F^1(A) \longrightarrow \dots$$

If (F^n) is cohomological then $F^{-1}(C) = 0$ and we get left exactness, if (F^n) is homological then $F^1(A) = 0$ and we get right exactness. \square

To emphasize the fact that a universal δ -functor is uniquely determined by F^0 we make the following definition:

Definition 5. *Given a universal cohomological δ -functor (F^n) we write $F^i = R^i F$, where $F = F^0$, and call $R^i F$ the **i -th right derived functor** of F . Given a universal homological δ -functor (F^n) we write $F^i = L^{-i} F$, where $F = F^0$, and call $L^i F$ the **i -th left derived functor** of F .*

Our goal is to show that under some mild conditions the necessary condition from Proposition 1 are also sufficient for a functor to be F^0 of some universal δ -functor.

Definition 6. *Let (F^n) be a universal cohomological (homological) δ -functor. An object A is called **acyclic** (for the δ -functor) if $F^n(A) = 0$ for all $n > 0$ (resp. all $n < 0$).*

We are now going to concentrate on cohomological functors but in parenthesis we will put remarks about analogous statements for homological δ -functors. Note that if $f : X \longrightarrow Y$ is a monomorphism (epimorphism) and Y is acyclic (X is acyclic) then f is an effacing monomorphism (epimorphism). Moreover, suppose that $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$ is a short exact sequence with B acyclic. In the long exact sequence

$$\dots \longrightarrow F^{k-1}(C) \longrightarrow F^k(A) \longrightarrow F^k(B) \longrightarrow F^k(C) \longrightarrow F^{k+1}(A) \longrightarrow \dots$$

we have $F^i(B) = 0$ for all $i > 0$ ($i < 0$ if the δ -functor is homological). It follows that the maps $\delta_k : F^k(C) \longrightarrow F^{k+1}(A)$ are isomorphisms for $k > 0$. This observation often allows to reduce statements about F^n to statements about F^{n-1} and is called **dimension shifting**. A more refined version of this ideas is the following fundamental

Theorem 2. *Let (F^n) be a cohomological universal δ -functor and set $F = F^0$. Given an exact sequence*

$$0 \longrightarrow A \xrightarrow{f} M_0 \xrightarrow{d_0} M_1 \xrightarrow{d_1} M_2 \xrightarrow{d_2} \dots$$

with M_i acyclic for all i , the k -th cohomology of the complex

$$(2) \quad 0 \longrightarrow F(M_0) \xrightarrow{F(d_0)} F(M_1) \xrightarrow{F(d_1)} F(M_2) \xrightarrow{F(d_2)} \dots$$

is isomorphic to $F^k(A)$ for all $k \geq 0$.

Similarly, let (F^n) be a homological universal δ -functor and set $F = F^0$. Given an exact sequence

$$0 \longleftarrow A \longleftarrow M_0 \longleftarrow M_1 \longleftarrow M_2 \longleftarrow \dots$$

with M_i acyclic for all i , the k -th cohomology of the complex

$$0 \longleftarrow F(M_0) \longleftarrow F(M_1) \longleftarrow F(M_2) \longleftarrow \dots$$

is isomorphic to $F^{-k}(A)$ for all $k \geq 0$.

Proof: We will give a proof for cohomological functors. We proceed by induction on k .

$k = 0$: Since the functor F is left exact, the sequence

$$0 \longrightarrow F(A) \longrightarrow F(M_0) \longrightarrow F(M_1)$$

is exact, so $F(A)$ is isomorphic to the kernel $F(d_0)$, i.e. to H^0 of the complex 2.

$k = 1$: Let A_1 be the image of d_0 (which is canonically isomorphic to the cokernel of f), so we have the following exact sequences:

$$0 \longrightarrow A \longrightarrow M_0 \xrightarrow{d} A_1 \longrightarrow 0$$

and

$$0 \longrightarrow A_1 \xrightarrow{f} M_1 \xrightarrow{d_1} M_2 \xrightarrow{d_2} M_3 \xrightarrow{d_3} \dots$$

The long exact sequence associated to the first of these sequences is

$$0 \longrightarrow F(A) \longrightarrow F(M_0) \longrightarrow F(A_1) \longrightarrow F^1(A) \longrightarrow F^1(M_0) = 0 \longrightarrow \dots$$

Thus $F^1(A)$ is a cokernel of the morphism $F(d) : F(M_0) \longrightarrow F(A_1)$. On the other hand, left exactness of F implies that the sequence

$$0 \longrightarrow F(A_1) \longrightarrow F(M_1) \longrightarrow F(M_2)$$

is exact, so $F(A_1)$ is a kernel of $F(d_1)$. Thus H^1 of the complex 2 is isomorphic to a cokernel of $F(d)$. Thus $F^1(A)$ and H^1 are canonically isomorphic.

inductive step: Suppose that the result holds for $k < n$, where $n \geq 2$.

The long exact sequence associated to

$$0 \longrightarrow A \longrightarrow M_0 \xrightarrow{d} A_1 \longrightarrow 0$$

shows that $F^n(A)$ is isomorphic to $F^{n-1}(A_1)$ (dimension shifting). Since the sequence

$$0 \longrightarrow A_1 \xrightarrow{f} M_1 \xrightarrow{d_1} M_2 \xrightarrow{d_2} M_3 \xrightarrow{d_3} \dots$$

is exact, we may use the inductive assumption and conclude that $F^{n-1}(A_1)$ is isomorphic to H^{n-1} of the complex

$$0 \longrightarrow F(M_1) \xrightarrow{F(d_1)} F(M_2) \xrightarrow{F(d_2)} F(M_3) \xrightarrow{F(d_3)} \dots$$

which is the same as H^n of the complex 2 (since $n \geq 2$). \square

Theorem 2 suggests a method to recover F^n from F^0 . The only problem is that we need to use acyclic objects, which are defined by using the functors F^n . Fortunately, we have the following crucial observation

Proposition 2. *Injective (projective) objects are acyclic for any universal cohomological (homological) δ -functor.*

Proof: We give a proof for cohomological δ -functors. Let I be an injective object and $u : I \longrightarrow Y$ an effacing monomorphism. The injectivity of I implies the existence of a morphism $w : Y \longrightarrow I$ such that $wu = id$. Thus, for $n > 0$, we have $id = F^n(id) = F^n(wu) = F^n(w)F^n(u) = F^n(w)0 = 0$, hence $F^n(I) = 0$ (an object is 0 iff $0 = id$). \square