## Homework 1

due on Wednesday, September 25

Study chapter 7 of Dummit and Foote. Solve problems 33 and 34 to 7.4.

**Problem 1.** An element a of a ring R is called **nilpotent** if  $a^m = 0$  for some m > 0.

- a) Prove that in a commutative ring R the set N of all nilpotent elements of R is an ideal. This ideal is called the **nilradical** of R. Prove that 0 is the only nilpotent element of R/N.
- b) Let R be a commutative ring and let  $a_1, ..., a_n \in R$  be nilpotent. Set I for the ideal  $\langle a_1, ..., a_n \rangle$  generated by  $a_1, ..., a_n$ . Prove that there is a positive integer N such that  $x_1x_2...x_N = 0$  for any  $x_1, ..., x_N$  in I (i.e. that  $I^N = 0$ ).
- c) Prove that the set of all nilpotent elements in the ring  $M_2(\mathbb{R})$  is not an ideal.
- d) Prove that if p is a prime and m > 0 then every element of  $\mathbb{Z}/p^m\mathbb{Z}$  is either nilpotent or invertible.
- e) Find the nilradical of  $\mathbb{Z}/36\mathbb{Z}$  (by correspondence theorem, it is equal to  $n\mathbb{Z}/36\mathbb{Z}$  for some n).

**Problem 2.** Let R be a commutative ring. For an ideal I of R define

$$\sqrt{I} = \{x \in R : x^n \in I \text{ for some } n > 0\}.$$

- a) Prove that  $\sqrt{I}$  is an ideal. It is called the **radical** of I.
- b) Prove that  $\sqrt{\{0\}}$  is the nilradical of R.
- c) Consider a surjective homomorphism  $f: R \longrightarrow S$ . Prove that in the correspondence theorem the nilradical of S corresponds to  $\sqrt{\ker f}$ .
- d) Prove that  $R/\sqrt{I}$  has trivial nilradical.
- **Problem 3.** A subset S of a commutative ring is called **multiplicative** if  $0 \notin S$  and for any  $a, b \in S$  also  $ab \in S$ .
- a) Let I be an ideal of a commutative unital ring R. Prove that I is a prime ideal iff R-I is multiplicative.

- b) Let S be a multiplicative subset of a comutative unital ring R. Consider the set T of all ideals of R which are disjoint with S. Prove that this set contains maximal elements (with respect to inclusion; this requires Zorn's Lemma and is very similar to the proof that every ring has a maximal ideal). Prove that every maximal element of T is a prime ideal.
- c) Use b) to prove that if  $a \in R$  is not nilpotent then there is a prime ideal in R which does not contain a.
- d) Prove that the nilradical of a commutative unital ring R coincides with the intersection of all prime ideals.

**Problem 4.** Let  $f: R \longrightarrow S$  be a homomorphism of commutative unital rings.

- a) Prove that if P is a prime ideal of S then  $f^{-1}(P)$  is a prime ideal of R. Is this true for non-commutative rings?
- b) Find an example when P is a maximal ideal of S but  $f^{-1}(P)$  is not maximal in R.
- c) Prove that if f is onto and Q is a prime ideal of R such that  $\ker f \subseteq Q$  then f(Q) is a prime ideal of S. Is this true for non-commutative rings?
- d) Suppose that f is surjective. Prove that if P is a maximal ideal of S then  $f^{-1}(P)$  is maximal in R. Prove that if Q is a maximal ideal of R then f(Q) is either S or it is a maximal ideal of S. Show by example that a similar statement for prime ideals is false.
- e) Find all prime ideals of  $\mathbb{Z}/36\mathbb{Z}$ .

**Problem 5.** Let R be a ring and n a positive integer. Prove that if I is an ideal of the ring  $M_n(R)$  then  $I = M_n(J)$  for some ideal J of R. Prove that I is maximal iff J is maximal. Prove that I is prime iff J is prime. Conclude that if R is simple (prime) then so is  $M_n(R)$ .

**Problem 6.** Let I be a prime ideal of R. Prove that if J, K are left ideals of R such that  $JK \subseteq I$  then either  $Jjj \subseteq I$  or  $K \subseteq I$ . Hint: Consider the set of all a such that  $aK \subseteq I$  and prove that it is an ideal containing J.