

NAME (Printed): _____ Section: _____

Math 304-5,6 Linear Algebra Fall 2018 Quiz 10 Feingold

INSTRUCTIONS: Show all calculations and reasons needed to justify your answers.

Let $V = \mathbf{R}^4$ with the standard dot product. Let $W = \langle T \rangle$ where

$$T = \left\{ w_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, w_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, w_3 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \right\}.$$

- (1) (4 Pts) Use the Gram-Schmidt process to convert T into an **orthogonal** basis $T' = \{w'_1, w'_2, w'_3\}$ for W .
-

(2) (3 Pts) Use your answer to part (1) to find the coefficients, x_1, x_2, x_3 , of the projection,

$Proj_W(v) = \sum_{i=1}^3 x_i w'_i$ of the vector $v = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in V$ onto the subspace W . They are

uniquely determined by the condition that $v - Proj_W(v)$ is orthogonal to W , that is, $(v - Proj_W(v)) \cdot w'_j = 0$ for $1 \leq j \leq 3$.

INSTRUCTIONS: Show all calculations and reasons needed to justify your answers.

Let $V = \mathbf{R}^4$ with the standard dot product. Let $W = \langle T \rangle$ where

$$T = \left\{ w_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, w_2 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, w_3 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \right\}.$$

- (1) (4 Pts) Use the Gram-Schmidt process to convert T into an **orthogonal** basis $T' = \{w'_1, w'_2, w'_3\}$ for W .

Solution: Step 1: $w'_1 = w_1$. Step 2: $w'_2 = w_2 - \frac{w_2 \cdot w'_1}{w'_1 \cdot w'_1} w'_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \frac{2}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}$.

Step 3: $w'_3 = w_3 - \frac{w_3 \cdot w'_1}{w'_1 \cdot w'_1} w'_1 - \frac{w_3 \cdot w'_2}{w'_2 \cdot w'_2} w'_2 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix} - \frac{4}{2} \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix} - \frac{6}{2} \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix}$.

So $T' = \left\{ w'_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, w'_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \\ 1 \end{bmatrix}, w'_3 = \begin{bmatrix} -1 \\ -1 \\ 1 \\ 1 \end{bmatrix} \right\}$.

Check that $w'_i \cdot w'_j = 0$ for $1 \leq i < j \leq 3$, and by the process, $\langle T' \rangle = \langle T \rangle$.

- (2) (3 Pts) Use your answer to part (1) to find the coefficients, x_1, x_2, x_3 , of the projection,

$Proj_W(v) = \sum_{i=1}^3 x_i w'_i$ of the vector $v = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \in V$ onto the subspace W . They are

uniquely determined by the condition that $v - Proj_W(v)$ is orthogonal to W , that is, $(v - Proj_W(v)) \cdot w'_j = 0$ for $1 \leq j \leq 3$.

Solution: The conditions mean that $v \cdot w'_j = Proj_W(v) \cdot w'_j = x_j (w'_j \cdot w'_j)$ for $1 \leq j \leq 3$ since T' is an orthogonal set. This says $x_j = \frac{v \cdot w'_j}{w'_j \cdot w'_j}$ so from part (1),

$$x_1 = \frac{a+c}{2}, \quad x_2 = \frac{b+d}{2}, \quad x_3 = \frac{-a-b+c+d}{4}.$$